


Anatomy of the Neck



- 1. Parotid gland
- 2. Superficial temporal artery & vein
- 3. Temporal branch of facial nerve
- 4. External carotid artery & posterior facial vein
- 5. Superficial cervical lymph nodes
- 6. External jugular vein
- 7. Accessory nerve & internal carotid artery
- 8. Platysma muscle
- 9. Fourth cervical nerve
- 10. Superior position of sternocleidomastoid muscle
- 11. Deep cervical lymph nodes
- 12. Fifth cervical nerve

- 13. Posterior supraclavicular nerve & anterior jugular vein
- 14. Superficial cervical artery & vein
- 15. Middle supraclavicular nerve & subclavian artery
- 16. Transverse scapular artery & vein
- 17. Inferior position of sternocleidomastoid muscle
- 18. External maxillary artery & anterior facial vein
- 19. Submaxillary lymph nodes & digastric muscle
- 20. Submaxillary gland & mylohyoid muscle
- 21. Submental lymph nodes & hypoglossal nerve
- 22. Superior laryngeal artery & nerve

- 23. Superior cervical ganglion
- 24. Superior laryngeal vein & omohyoid muscle
- 25. Superior thyroid artery & vein
- 26. Ansa hypoglossi
- 27. Common carotid artery & sternothyroid muscle
- 28. Middle cervical ganglion & phrenic nerve
- 29. Vagus nerve
- 30. Thyroid gland & middle thyroid vein
- 31. Internal jugular vein
- 32. Sternohyoid muscle
- 33. Jugular lymphatic trunk
- 34. Inferior thyroid veins

This is one of a series of paintings for Lederle by Paul Peck, illustrating the anatomy of various organs and tissues of the body which are frequently attacked by infection, where aureomycin may prove useful.





is valuable in

Infections of the Neck

and is especially

useful where surgery

is indicated

Literature available on request-

LEDERLE LABORATORIES DIVISION

AMERICAN Cyanamid company
30 ROCKEFELLER PLAZA, NEW YORK 20, N. Y.

for peptic ulcer patients 8 hours' relief from a single dose

PRANTAL RA

TABLETS

first repeat action anticholinergic



less frequent dosage uninterrupted night rest greater freedom from side effects

PRANTAL*
3 forms
for more
flexible therapy

PRANTAL Repeat Action Tablets, 100 mg. Dosage: One or two tablets every eight hours. PRANTAL Tablets (plain), 100 mg., scored. Dosage: One or two tablets every six hours. PRANTAL Injection (subcutaneous or intramuscular), 25 mg. per cc., 10 cc. vials. Dosage: 0.5 mg. per Kg. of body weight every six hours.

*T. M. Brand of diphenmethanil methylsulfate

Schering corporation · bloomfield, new jersey



Announcing



Gantricillin is the new combination of Gantrisin 'Roche' (the single, more soluble sulfonamide) plus penicillin.

Gantricillin is recommended for infections susceptible to penicillin or sulfonamides. It is especially useful when the causative organisms are more susceptible to the combination than to either drug alone. Each scored tablet contains 0.5 Gm Gantrisin and 100,000 units of crystalline penicillin G potassium.

Hoffmann-La Roche Inc., Nutley 10, N. J.

GANTRISIN®—brand of sulfisoxazole GANTRICILLIN™

A complete lipotropic formula in high potency dosage

HEPATRÓPIN

TRADE MARK

LIPOTROPIC MIXTURE U.M.A.

For the treatment of

LIVER DISORDERS

as they occur in

CIRRHOSIS ATHEROSCLEROSIS ALCOHOLISM DIABETES

and where there is

IMPAIRED METABOLISM OF FAT AND CHOLESTEROL

4 tablets t.i.d. supplies		
Choline bitartrate	5.0 Gm.	HEPATROPIN TABLETS
dl. Methionine	0.6 Gm.	are supplied in
Inositol	1.0 Gm.	Bottles of 100
Folic acid	1.5 mg	boilles of 100
Vitamin B ₁₂	25.0 mcgm	Literature on request



PHARMACEUTICAL DIVISION

New York 3, N. Y.

PROVEN PAIN CONTROL

with safety

'Tabloid'

'EMPIRIN' COMPOUND

with CODEINE PHOSPHATE*


gr. $\frac{1}{8}$ No. 1 gr. $\frac{1}{4}$ No. 2 gr. $\frac{1}{2}$ No. 3 gr. 1 No. 4

*12 times more soluble than sulfate



Burroughs Wellcome & Co. (U.S.A.) Inc. Tuckahoe 7, N. Y.

ETHICON SURGICAL SUTURES



to prevent attacks in angina pectoris

Nocturnal angina pectoris (as well as day-time attacks) responds gratifyingly to Peritrate. Investigators report that in patients on Peritrate "insomnia due to pain was much reduced and they were able to enjoy reasonably good sleep." Others had less palpitation during the night "and were able to sleep on the left side" which they had been unable to do prior to Peritrate. When required, an extra 1-1½ tablets taken on retiring brought "striking relief."

Prophylactic Management

Peritrate—a long-lasting coronary vasodilator—prevents rather than relieves attacks. The prophylactic action of each dose lasts 4 to 5 hours. One to 2 tablets of Peritrate, taken 3 to 4 times daily, can...

- 1. reduce the number of attacks and
- 2. reduce the severity of non-preventable attacks.

Effective in 4 out of 5 Patients

Fewer or less severe attacks occurred in 80% and 78.4% of patients, 1,2 and in the majority the need for nitroglycerin was reduced. Clinical improvement has been verified by EKG findings and exercise tolerance may improve measurably. 2,3

Available in 10 mg. tablets in bottles of 100, 500 and 5000.

Literature and samples on request.

- 1. Humphreys, P., et al.: Angiology 3:1 (Feb.) 1952.
- 2. Plotz, M.: New York State J. Med. 52:2012 (Aug. 15) 1952.
- 3. Perlman, A.: Angiology 3:16 (Feb.) 1952.



WARNER-CHILCOTT


Laboratories NEW YORK

76.6% of patients preferred

EURAX[®]

for relief of





In this study, as in previous reports²⁻⁵, EURAX Cream produced complete relief of itching in approximately 65 per cent of cases, and partial relief in most of the remainder.

Other favorable features of EURAX Cream that were again confirmed include:

- /, Prolonged effect lasting up to 8 hours or more.
- √, No loss of effect on continued use.
- ✓ Virtually complete lack of sensitizing or toxic properties.


EURAX... not an antihistaminic or a -caine derivative... is indicated for prompt, prolonged relief of itch in practically all forms of dermatosis including pruritus due to administration of antibiotics.

EURAX Cream* (brand of crotamiton cream) contains 10% N-ethyl-o-crotonotoluide in a vanishing-cream base. Tubes of 20 Gm. and 60 Gm. and jars of 1 lb. at your local pharmacy.

Bibliography

1. Hitch, J. M.: North Carolina M. J. 12:548, 1951. 2. Peck, S. M., and Michelfelder, T. J.: New York State J. Med. 50:1934, 1950. 3. Couperus, M.: J. Invest. Dermat. 13:35, 1949. 4. Soifer, A.: Quart. Rev. Int. Med. & Dermat. 8:1, 1951. 5. Johnson, S. M., and Bringe, J. W.: Arch. Dermat. & Syph. 63:768, 1951. *U.S. Pat. #2,505,681.

Samples and Reprints on Request



New, Improved 'Homicebrin' with Vitamin B₁₂

'Homicebrin' now provides <u>six</u> essential vitamins in a smooth, palatable, homogenized mixture.

Each 5 cc. contain:

Vitamin A (Palmitate)	3,000 U.S.P	. units
Thiamin Chloride		mg.
Riboflavin	: 1.:	2 mg.
Vitamin B ₁₂ (Activity Equiv	valent) 3	mcg.
Ascorbic Acid	60	mg.
Vitamin D	1.000 U.S.P	. units

Prescribe the 4-ounce or the economical 1-pint bottle. There is no increase in price.

Eli Lilly and Company Indianapolis 6, Indiana, U.S.A.





For the prophylaxis or treatment of multiple vitamin deficiencies in infants and children.

Homicebrin

(HOMOGENIZED MULTIPLE VITAMINS, LILLY)

Searle Research Progress Report:

Continued investigational work has resulted in Pro-Banthine, a new anticholinergic drug with high potency, small dosage, minimal side effects, agreeable taste and convenient dosage schedule.

The new anticholinergic, Pro-Banthine* (brand of propantheline bromide) provides a powerful drug in the therapy of peptic ulcer, intestinal hypermotility and other conditions of parasympathotonia.

The high potency of Pro-Banthine permits its use in small dosage. With the suggested dosage of one tablet (15 mg.) with meals and two at bedtime there is little likelihood of untoward manifestations.

Pro-Banthine has a pronounced inhibiting action on stimuli at (a) the parasympathetic and sympathetic ganglia and (b) the effector organs of the parasympathetic system.

Pro-Banthine is produced for oral use in 15 mg. sugar-coated tablets.


SEARLE


Research in the Service of Medicine *Trademark of G. D. Searle & Co.


Top—Section through duodenal bulb just distal to pylorus through center of ulcer crater.


Center—Healing ulcer with scar tissue and regeneration of tissue layers.

Bottom-Healed ulcer with restoration of mucosa.









<u>Levugen</u>

fructose
for intravenous
infusion...
now, for the
first time,
available
for general
clinical

use

(FRUCTOSE, MEAD)

The advantages of fructose for intravenous infusion have long been recognized. But limited availability of pure fructose has prevented its general clinical use. Now, extensive research in carbohydrate chemistry by Mead Johnson & Company has resulted in a practical and economical method of producing pure fructose. As Levugen 10% in Water, it is available for intravenous use.

Levugen (Fructose, Mead) can be infused much more rapidly than dextrose, with better retention and less disturbance of fluid balance.

Since Levugen is removed from the blood very rapidly, it does not produce high hyperglycemic levels or spill over into the urine in significant amounts even when it is infused in fairly high concentration. Levugen can therefore be given much more rapidly than dextrose, with less loss of calories through glycosuria. A liter of Levugen 10% in Water can be given in the same time as a liter of 5% dextrose.

Clinical Advantages of Levugen

- More rapid provision of calories
- Less loss of calories through glycosuria
- More rapid formation of liver glycogen
- Less disturbance of fluid balance
- Shorter infusion time, with less discomfort for the patient
- Less time and trouble for hospital personnel

"<u>Levugen</u>



can be infused more rapidly"

Levugen 10% in Water is available in 1 liter (1000 cc.) flasks, containing 100 Gm. of Levugen (Fructose, Mead), approximately 400 calories.



Levugen

MEAD JOHNSON & COMPANY Evansville 21, Ind., U.S.A. MEAD



CAMBRIDGE MOBILE MODEL



"SIMPLI-TROL"
PORTABLE

"SIMPLI-SCRIBE" DIRECT WRITER



"CAMBRIDGE has it!"



EDUCATIONAL CARDIOSCOPE



OPERATING ROOM CARDIOSCOPE

The first Cambridge Electrocardiographs were made available to the medical profession over thirty-five years ago. It is fortunate that from the beginning, these instruments produced consistently accurate electrocardiograms, since it was upon these records that the science of Electrocardiography has been largely built in this country. It follows, therefore, that the Medical Profession looks to CAMBRIDGE for new instruments now demanded in the tremendous development the science is undergoing.

Today, CAMBRIDGE MOBILE and "SIMPLI-TROL" MODEL ELECTROCARDIOGRAPHS are the standard of comparison. CAMBRIDGE RESEARCH MODEL meets the most exacting demands of advanced cardiac study. CAMBRIDGE "SIMPLI-SCRIBE" DIRECT WRITER gives the doctor or institution a fine, convenient, supplementary electrocardiograph.

CAMBRIDGE EDUCATIONAL CARDIOSCOPE is the newest aid in teaching Electrocardiography and auscultation by visual demonstration. The CAMBRIDGE OPERATING ROOM CARDIOSCOPE provides a continuous tracing of any lead of the electrocardiogram, indicates the instantaneous heart rate and warns of approaching cardiac standstill during surgery. In the opinion of qualified anesthesiologists and surgeons, this instrument will soon be a "must" for the modern operating room.

Send for descriptive literature

CAMBRIDGE INSTRUMENT CO., INC. 3744 Grand Central Terminal, New York 17, N.Y.

Cleveland 15, 1720 Euclid Avenue Philadelphia 4, 135 South 36th Street Chicago 12, 2400 West Madison Street

PIONEER MANUFACTURERS OF THE ELECTROCARDIOGRAPH

CAMBRIDGE ALSO MAKES ELECTROKYMOGRAPHS, PLETHYSMOGRAPHS, AMPLIFYING STETHOSCOPES, RESEARCH pH METERS, BLOOD PRESSURE RECORDERS, INSTRUMENTS FOR MEASURING RADIOACTIVITY, ETC.



Herald Square, N. Y.

Parkchester, Bronx



Jamaica, Queens

Flatbush, Brooklyn


White Plains, N. Y

MACY'S PRESCRIPTION DEPARTMENT

Macy's cooperates with the medical profession to control the indiscriminate use of dangerous drugs. If a patient wishes to refill a prescription containing any drug which may be harmful if used without supervision, the physician is immediately notified. And with the physician's approval, the prescription is accurately compounded, carefully checked.

No wonder so many physicians prefer Macy's Prescription Department.

......





FULL SPEED AHEAD in TISSUE REPAIR

DESITIN Ointment proves in everyday practice its ability to ease pain, renew vitality of sluggish cells, and stimulate smooth tissue repair in lacerated, denuded, chafed, irritated, ulcerated tissues—in conditions often resistant to

other therapy. 1-3

OINTMENT
the pioneer external cod liver oil therapy
in WOUNDS (especially slow healing)

in Wounds (especially slow healing)
ulcers (decubitus, varicose, diabetic)
burns, perianal dermatitis
non-specific dermatoses

Protective, soothing, healing, Desitin Ointment is a non-irritating blend of high grade, crude Norwegian cod liver oil (with its unsaturated fatty acids and high potency vitamins A and D in proper ratio for maximum efficacy), zinc oxide, talcum, petrolatum, and lanolin. Desitin Ointment does not liquefy at body temperature and is not decomposed or washed away by secretions, exudate, urine or excrements. Dressings easily applied and painlessly removed.

Tubes of 1 oz., 2 oz., 4 oz., and 1 lb. jars.

write for samples and literature

DESITIN CHEMICAL COMPANY

70 Ship Street • Providence 2, R. I.

- Behrman, H. T., Combes, F. C., Bobroff, A., Leviticus, R.: Ind. Med. & Surg. 18:512, 1949.
- 2. Turell, R.: New York St. J. M. 50:2282, 1950.
- 3. Heimer, C. B., Grayzel, H. G., and Kramer, B.: Archives Pediat. 68:382, 1951.

FIFTEENTH SERIES

POSTGRADUATE RADIO PROGRAM

THE NEW YORK ACADEMY OF MEDICINE

Committee on Medical Information in cooperation with Committee on Medical Education

2 East 103 Street, New York

Thursdays -9:00 p.m. -10:00 p.m. . . . Station WNYC-FM -93.9 megs.

APRIL 2

TREATMENT OF DIABETIC KE-TOSIS AND COMA

Edward Tolstoi, Associate Professor of Clinical Medicine, Cornell University Medical College.

THE MANAGEMENT OF THE EN-LARGED OVARY

M. A. Goldberger, Attending Gynecologist to The Mount Sinai Hospital.

APRIL 9

THE MEDICAL MANAGEMENT OF ACUTE AND CHRONIC ARTERIAL OCCLUSION

A. Wilbur Duryee, Professor of Clinical Medicine, New York University Post-Graduate Medical School.

SKIN CANCER

Anthony C. Cipollaro, Professor and Director, Department of Dermatology, New York Polyclinic Medical School and Hospital.

APRIL 16

HALSTED, MAN AND SURGEON Bertram M. Bernheim, Associate Professor of Surgery, The Johns Hopkins Medical School.

WILLIAM H. HALSTED; THE SUR-GEON'S PRECEPTOR

John J. Moorhead, Professor Emeritus of Surgery, New York University Post-Graduate Medical School.

CANCER OF THE BREAST

William C. White, Clinical Professor of Surgery, Columbia Universiy, College of Physicians and Surgeons.

APRIL 23

METABOLIC EFFECTS OF INSULIN DeWitt Stetten, Jr., Chief, Division of Nutrition and Physiology, The Public Health Research Institute of the City of New York.

PRECANCEROUS LESIONS OF THE **VULVA**

Mortimer D. Speiser, Associate Clinical Professor of Obstetrics and Gynecology, New York University Medical College.

APRIL 30

PAUL EHRLICH, MAN AND SCIEN-TIST


Selman A. Waksman, Professor of Microbiology, Rutgers University, New Bruns-wick, N. J.

CHEMOTHERAPY PAUL EHRLICH

Perrin H. Long, Professor of Medicine and Chairman of the Department of Medicine, College of Medicine, State University Medical Center at N.Y.C.

USE OF HORMONES IN THE PAL-LIATIVE TREATMENT OF MAM-MARY CARCINOMA

George C. Escher, Assistant, Memorial Hospital, Section Head, Hormone Chemotherapy.



Upjohn

mixed surface infections...


Each gram contains 5 mg. neomycin sulfate (equivalent to 3.5 mg. neomycin base).

Available: Ointment in ½ oz. and 1 oz. tubes, and 4 oz. jars. Cream in ½ oz. tubes.

The Upjohn Company, Kalamazoo, Michigan



Trademark V. S. Pat. Off. Successful Success



PLASTIC AND GLASS EYES



MAGER & GOUGELMAN, Inc.

OVER A CENTURY OF SERVICE • 1851-1952

- PRIVATE FITTINGS
- PERSONAL ATTENTION
- EXPERIENCE & SKILL

510 MADISON AVE., N. Y. 22 Telephone PLaza 5-3756

functional, fashionable, full - footed .

NO OVERHOSE NECESSARY

with sheer ACE Elastic Hosiery

(fashioned by B-D, makers of ACE elastic bandage)

From the viewpoint of Function (important to you) and from the viewpoint of Fashion (important to patients), ACE Elastic Hosiery is a vastly superior product . . . as distinctive in its field as ACE Elastic Bandage is in its field. Here then is elastic hosiery your women patients cannot object to. Prescribe a pair and see how they will please the patient. Also available, ACE Elastic Hosiery—styled for men.

Available now at your "Professional Source of Supply"

—Gladys Gray's staff of Registered Nurses assures proper fitting and instructions.


Literature on request.



CAMP SUPPORTS . KENDRICK and BAUER & BLACK ELASTIC HOSIERY . "IDENTICAL" BREAST FORMS.

R. N. 133 EAST FIFTY-EIGHTH STREET NEW YORK 22, N. Y. SUITE 201

PLAZA 5-6328



CHARLES B. TOWNS HOSPITAL Established 1901

FOR THE TREATMENT OF ALCOHOLISM

NARCOTIC AND BARBITURATE

ADDICTIONS EXCLUSIVELY

Complete Medical & Psychiatric Treatment at predetermined cost. Privacy of patient is assured—if desired.

Literature on Request

Edward B. Towns, Director

293 Central Park West, N. Y. 24, N. Y. SChuyler 4-0770

Member American Hospital Association

Aminophyllin . . .

a "most effective single agent for prompt relief" of severe bronchial asthma

"useful as a peripheral vasodilator and myocardial stimulant" in

pulmonary edema paroxysmal dyspnea of congestive heart failure Cheyne-Stokes respiration



H. E. DUBIN LABORATORIES, INC. 250 E. 43rd St. • New York 17, N.Y.

dubin

aminophyllin

(theophylline-ethylenediamine)

readily soluble for rapid therapeutic effect.

TABLETS • AMPULS POWDER SUPPOSITORIES



Satisfaction Guaranteed Specialists in Plastic and Glass Artificial Human Eyes Exclusively

REFERRED CASES CAREFULLY ATTENDED

FRIED & KOHLER, Inc.

665 FIFTH AVENUE near 53rd Street

NEW YORK, N. Y. Tel. ELdorado 5-1970



in the hands of the physician

Often the critical evaluation of the drug to be administered is as important to the patient's recovery as is the diagnosis of his condition. In each case correct procedures can be determined only by the physician.

CHLOROMYCETIN is eminent among drugs at the disposal of the medical profession. Clinical findings attest that, in the hands of the physician, this widely used, broad spectrum antibiotic has proved invaluable against a great variety of infectious disorders.



notably effective well tolerated broad spectrum antibiotic

The many hundreds of clinical reports on CHLOROMYCETIN emphasize repeatedly its exceptional tolerance as demonstrated by the infrequent occurrence of even mild signs and symptoms of gastrointestinal distress and other side effects in patients receiving the drug.

Similarly, the broad clinical effectiveness of CHLOROMYCETIN has been established, and serious blood disorders following its use are rare. However, it is a potent therapeutic agent, and should not be used indiscriminately or for minor infections-and, as with certain other drugs, adequate blood studies should be made when the patient requires prolonged or intermittent therapy.

CHLOROMYCETIN (chloramphenicol, Parke-Davis) is available in a variety of forms, including: CHLOROMYCETIN Kapseals,® 250 mg., bottles of 16 and 100.

CHLOROMYCETIN Capsules, 100 mg., bottles of 25 and 100.

CHLOROMYCETIN Capsules, 50 mg., bottles of 25 and 100.

CHLOROMYCETIN Ophthalmic Ointment, 1%, 1/8-ounce collapsible tubes.

CHLOROMYCETIN Ophthalmic, 25 mg. dry powder for solution, individual vials with droppers.



available on prescription only

Quadrinal tablets

DRUG..

PARTICULAR ..

ASTHMA AND RELATE..

RESPIRATORY CONDITIONS.

Quadrinal Tablet

---demarks E QUADRINAL TABLETS CONTAIN FOUR DRUGS, EACH SELECTED FOR ITS PARTICULAR EFFECT IN CHRONIC **ASTHMA AND RELATED ALLERGIC**


1/2 or 1 Quadrinal Tablet every 3 or 4 hours, not more than three tablets a day.

Each Quadrinal Tablet contains ephedrine hydrochloride % gr. (24 mg.), phenobarbital % gr. (24 mg.), Phyllicin (theophylline-calcium salicylate) 2 gr. (120 mg.), and potassium iodide 5 gr. (0.3 Gm.)

Quadrinal Tablets are marketed in bottles of 100, 500 and 1000.

Quadrinal, Phyllicin, Trademarks E. Bilhuber, Inc.

distributor: BILHUBER-KNOLL CORP., Orange, New Jersey, U. S. A.



FOR Epileptic controlgrand mal and petit mal New Mebaroin

Highly Effective • Well Tolerated • Relatively Tasteless

In a comparative study of 108 patients^{1,2} with idiopathic epilepsy, by far the best results (85.6 per cent controlled or improved) were obtained with a combination of Mebaral and diphenylhydantoin. This combination achieved the closest approach to the ideal of maximum control of seizures with minimal toxicity. Each tablet of Mebaroin contains 90 mg. Mebaral® and 60 mg. diphenylhydantoin.

Dosage: Average adult dose: 1 or 2 tablets three times daily. For children over 6 years, 2 or 3 tablets daily; younger children ½ tablet once or twice daily. Treatment is best begun with a small dose which is increased gradually until the individual optimum is established. The usual caution should be observed when changing from another antiepileptic drug to Mebaroin.

Supplied: Bottles of 100 and 1000 scored tablets.

Meller, R. L., and Resch, J. A.: Bull. Univ. Minnesota Hosp., 20:78, Oct. 8, 1948.

Meller, R. L., and Resch, J. A.: Postgrad. Med., 6:452, Dec., 1949.



baby needs. LOW FAT

the premature

HIGH PROTEIN

ipso facto.

DRYCO°

"a half-skimmed milk with Vitamin A and D fortification"

In the report of a study of 122 premature infants whose birth weights were between 1,000 and 2,000 grams, Gordon* states, "There can be no doubt that under conditions of modern hospital practice, with cleanliness, good refrigeration and careful supervision of feeding, the use of a half-skimmed cow's milk mixture will produce more satisfactory weight gains [than evaporated milk or human milk]."

Prescribe Dryco for your "prematures"...and for other feeding problems. Write for your copy of suggested Dryco feeding instructions.

*Gordon, H. H.: Feeding of Premature Infants, Am. J. Dis. Child. June 1947.

DRYCO COSTS LESS PER QUART GENERALLY THAN GRADE A PASTEURIZED WHOLE MILK


DRYCO—a successful infant food for more than thirty years







THE BORDEN COMPANY 350 Madison Ave., New York 17 **Prescription Products Division**



$free\ flow$

IN URINARY TRACT INFECTIONS

Free flow of urine is essential to successful chemotherapy. Thus, it is important to select the sulfonamide preparation which is least likely to produce crystalluria.

On the score of both potency and solubility in acid urine, SULFOSE is unsurpassed.

SULFOSE


Triple Sulfonamides Wyeth

ORAL SUSPENSION

TABLETS



Philadelphia 2, Pa.



An Important Diet-Feature in Weight-Control Programs

WALKER-GORDON CERTIFIED LOW-FAT Skimmed MILK

Remove the cream from Walker-Gordon Certified Whole Milk and you have Walker-Gordon Certified Skimmed Milk... custom-made for weight-control and weight-reduction programs. Your patients get the necessary nutrients of Walker-Gordon Certified Whole Milk without the butter fat and fat soluble vitamins. All the minerals including calcium and phosphorus, water-soluble vitamins, amino acids, and proteins remain, but only about half the calories of Whole Milk are present. And this pure, delicious, fresh Skimmed Milk has real taste-appeal.

The Medical Profession also frequently recommends Walker-Gordon Certified Skimmed Milk in cases of Pregnancy and Lactation, Childhood and Adult Obesity, Abnormal Bile Secretion, Celiac Disease, Infant Feeding, Gastric Ulcers, Diarrhea, Psoriasis, Allergy, Diabetes, Colitis, Acne, Eczema, and Hypertension.

Delivered fresh within one day of milking by leading dairy distributors in New York, New Jersey, and Pennsylvania.


FREE! Descriptive book, "Technical Control and Supervision of Walker-Gordon Certified Milk," sent without obligation on request.

Walker-Gordon Laboratory Co.

Plainsboro, N. J.

Phone Plainsboro, 3-2750


Certified by the Medical Milk Commissions of the Counties of New York, Kings, Hudson, and Philadelphia





IN URINARY

TRACT INFECTIONS



"Patients with pyelitis were well and doing their usual duties within 24 hours . . . "1 ". . . resistant cases showed remarkable response."2

high urine levels

"Terramycin was selected . . . in view of high urinary excretion rate following small oral doses of the antibiotic."1

unexcelled toleration

"Terramycin is generally well tolerated, the percentage of relapses being low and the percentage of bacteriological as well as clinical cures high."3

- 1. Canad. M. A. J. 66:151 (Feb.) 1952.
- 2. J. Urol, 67:762 (May) 1952.
- 3, Ibid. 69:315 (Feb.) 1953.

erramycin of systemacycline

oral diuretic without equal

- "... superior... in promoting sodium and water excretion."1
- "... three-fourths the diuretic action of the standard [meralluride by injection]..."2
- "...a valuable substance to replace parenteral diuretics in patients who require continuous diuretic medication."3

NEOHYDRIN

THE DIURETIC TABLETS THAT WORK



- Moyer, J. H., and Handley, C. A.: Federation Proc. 11:378, 1952.
 Greiner, T.; Gold, H.; Warnhaw, L.; Palumbo, F.; Weaver, J.; Mathes, S., and Marsh, R.: Federation Proc. 11:352, 1952.
 Goldman, B. R., and Steigmann, F.: J. Lab. & Clin. Med. 40:803, 1952.

how to use this new drug

Maintenance of the edema-free state has been accomplished with as little as one or two NEOHYDRIN Tablets a day. Often this dosage of NEOHYDRIN will obtain per week an effect comparable to a weekly injection of MERCUHYDRIN.® When more intensive therapy is required one or two tablets three times daily may be prescribed as determined by the physician.

Gradual attainment of intensive therapy is recommended to preclude gastrointestinal upset which may occur in occasional patients with immediate high dosage. In rare instances a sensitivity to NEOHYDRIN may arise. Though sustained, the onset of NEOHYDRIN diuresis is gradual. Injections of MERCUHYDRIN will be initially necessary in acute severe decompensation. Contraindicated in acute nephritis and nephrosclerosis. Any patient receiving a diuretic should ingest daily a glass of orange juice or other supplementary source of potassium. Any patient receiving a diuretic should be watched for signs of depletion in sodium and chlorides especially in hot weather. Such depletion may first manifest itself as a refractivity to the diuretic and can be corrected by ingestion of sodium chloride.

Bottles of 50 tablets. There are 18.3 mg. of 3-chloromercuri-2methoxy-propylurea in each tablet.





... for the more common bacterial infectious diseases



Just 1 or 2 Pentids Tablets t.i.d. are particularly effective ... convenient, easy-to-take ... cause fewer side effects ... and are less than $\frac{1}{2}$ the cost of the newer antibiotics.

Bottles of 12 and 100.

formulated for convenient t.i.d. dosage

Pentids
Squibb 200,000 Unit Penicillin Tablets